



This preprint version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/> Restricted use of this manuscript is permitted provided the original work is properly cited. The authors assert their moral rights, including the right to be identified as an author.



Liquid Chromatographic Method for the Determination of Lutein in Milk and Pediatric Formulas

Brendon D. Gill*, and Harvey E. Indyk

Fonterra Waitoa, PO Box 7, Waitoa, New Zealand

* Corresponding author

Abstract

A simple and rapid method has been developed for routine compliance testing of lutein in pediatric formulas. Following a mild saponification procedure (70 °C for 10 min) demonstrated to be benign to lutein, lipophilic components were partitioned by a single extraction into hexane:diisopropyl ether (75:25 v/v) and concentrated in mobile phase. Baseline chromatographic separation of lutein and zeaxanthin (internal standard) was achieved using a C₃₀ column and isocratic elution with methanol:dichloromethane (70:30 v/v). Lutein concentration was quantitated against zeaxanthin, which corrected for incomplete extraction recovery. Performance parameters assessed included recovery (101–108%) and repeatability (2.2% RSD). The method was applied to the analysis of lutein supplemented pediatric formulas, unsupplemented milk powders, and bovine and human milk.

1. Introduction

Carotenoids are a diverse class of polyisoprenoid hydrocarbons (carotenes) and their oxygenated derivatives (xanthophylls), of which approximately 600 have been isolated from natural sources, predominantly in their most thermodynamically stable all-trans configuration. Within this group, there is a wide range of polarity, solubility and stability, and these features are of significance in the development of suitable analytical methods for their quantitative estimation in foods (Craft & Soares, 1992).

Carotenoid structure confers remarkably diverse physiological functions in photosynthesis, pro-vitamin A activity, antioxidant status, immune function, cell differentiation and several pathologies (Oliver & Palou, 2000). Since lutein and zeaxanthin are the only dietary carotenoids in the macular pigment, they have received increasing attention for their dietary potential to reduce age-related macular degeneration (Lietz, Mulokozi, Henry, & Tomkins, 2006), although this view has recently been challenged (Trumbo & Ellwood, 2006). Further, lutein and zeaxanthin are both considered to be protective in neonatal retinal epithelium, and infant formulas are therefore increasingly supplemented with lutein to more closely resemble human breast milk (Jewell, Mayes, Tubman, Northrop-Clewes, & Thurnham, 2004).

Contemporary methods for the selective analysis of carotenoids in foods and clinical applications are based on high performance liquid chromatography (HPLC), and comprehensive recent reviews are available (Barua, Olson, Furr, & van Breemen, 2000; Blake, 2007; Eitenmiller & Landen, 1999; Oliver & Palou, 2000; Su, Rowley, & Balazs, 2002). Photodiode-array detection is considered to be essential, given the subtle differences in characteristic UV–vis spectra of the many individual carotenoids potentially present in natural foods, while mass spectrometric detection methods have also been advocated for confirmation of identity (Breithaupt, 2004; Dachtler, Glaser, Kohler, & Albert, 2001; Rodríguez-Bernaldo de Quirós & Costa, 2006; Schlatterer & Breithaupt, 2006). Reversed phase separation systems are most commonly described utilizing monomeric or polymeric C₁₈ chemistries, although C₃₀ columns are increasingly applied in view of improved selectivity (Aman, Bayha, Carle & Schieber, 2004; Breithaupt, 2004; Dachtler et al., 2001; Li, Qin, Yin, & Tang, 2007; Mercadante, Steck, & Pfander, 1999; Rousell, Raley, & Hofsommer, 1996). Nevertheless, normal phase separations have also been reported in view of their enhanced retention of polar xanthophylls and resolution of isomeric carotenoids (Panfili, Fratianni, & Irano, 2004).

Saponification and direct solvent extraction methods are generally advocated for carotenoid analysis of foods, with the latter technique often preferred for low fat samples and when efficient extraction of target analytes can be demonstrated. However, for complex foods and those containing high fat,

alkaline hydrolysis is frequently required to remove potentially interfering lipid and chlorophyll, improve recovery and convert xanthophyll esters (Ball, 2006; Barua et al., 2000; Eitenmiller & Landen, 1999; Hart & Scott, 1995; Konings & Roomans, 1997; Larsen & Christensen, 2005; Rodríguez-Bernaldo de Quirós & Costa, 2006). Supercritical fluid extraction and enzymatic lipolysis have also been described as alternatives to these conventional techniques (Lietz & Henry, 1997; Oliver & Palou, 2000).

The carotenoid profile of milk has been reviewed, and while human milk reportedly contains significant levels of lutein, cryptoxanthin, lycopene and β -carotene, the major forms in bovine milk are lutein and β -carotene (Calderón et al., 2007; Canfield, Giuliano, & Graver, 1995; Canfield et al., 2003; Jensen, 1995). Two recent reviews have described carotenoid metabolism and factors affecting expression in bovine milk (Nozière et al., 2006; Sauvant, Grolier, & Azaïs-Braesco, 2002). In view of their high fat content, saponification has generally been applied to the extraction of carotenoids in milk and infant formulas (Giuliano, Neilson, Kelly, & Canfield, 1992; Granelli & Helmersson, 1996; Indyk, 1987; Jackson, Lien, White, Bruns & Kuhlman, 1998; Jewell et al., 2004; Schweigert, Hurtienne & Bathe, 2000), although enzymatic digestion has also been described (Khachik et al., 1997; Liu, Xu, & Canfield, 1998).

This study describes a simple, robust and validated method for the determination of lutein in bovine and human milk, milk powders, and infant formulas involving a single-tube alkaline hydrolysis and solvent extraction scheme with internal standard quantitation.

2. Experimental

2.1. Apparatus

The HPLC system consisted of a SCL-10Avp system controller, LC-10ADvp pump, FCV-10ALvp low pressure gradient unit, SIL-10AF sample injector unit equipped with a 50 mL injection loop, DGU-14A degasser unit, CTO-10ASvp column oven and a SPD-M10Avp photodiode array detector (Shimadzu, Kyoto, Japan). Instrument control and data processing were implemented using Shimadzu Class-VP version 6.12.

Absorbances for calibration standards were acquired with a model UV-1601 spectrophotometer (Shimadzu) with digital readout to 4 decimal places. Samples were centrifuged in a Funke Gerber Super Vario-N (Berlin, Germany). Terumo 3 mL disposable syringes (Terumo Corporation, Laguna, Philippines) and Minisart 0.2 μ m syringe filters with cellulose acetate membranes (Sartorius, Göttingen, Germany) were used for standard preparation. Prior to use, mobile phases were filtered

and degassed using a filtration apparatus with 0.45 µm nylon filter membranes (All-Tech, Deerfield, IL, USA).

2.2. Reagents

Lutein and zeaxanthin were purchased from ChromaDex (Irvine, CA, USA). Potassium hydroxide, pyrogallol and HPLC grade solvents, hexane, diisopropyl ether, ethyl acetate, methanol, dichloromethane, and ethanol, were supplied by Merck (Darmstadt, Germany). Water was purified with resistivity >18 MΩ using an E-pure water system (Barnstead, Dubuque, IA, USA).

2.3. Standards

Due to the light-sensitive nature of carotenoids, all steps in this procedure were performed under subdued lighting. Stock standards were prepared by weighing approximately 1 mg of lutein and zeaxanthin into separate 25 mL volumetric flasks and dissolved in approximately 20 mL of ethanol accomplished by shaking and ultrasonication for 5 min. The stock standards were then made to volume with ethanol. Intermediate standards were prepared by filtering approximately 3 mL of the stock solution through a 0.22 µm syringe filter into a 100 mL volumetric flask. Approximately 80 mL of ethanol was added and the flask sonicated for 5 min, then made to volume with ethanol.

The concentration of individual intermediate standards was determined from their measured absorbance against ethanol and published absorptivities ($E^{1\%}$: lutein, 2550 at 445 nm and zeaxanthin, 2480 at 452 nm) (Scott, Finglas, Seale, Hart, & de Froimont-Gortz, 1996) and corrected for chromatographic purity.

Five calibration standards were prepared containing a constant concentration of the internal standard zeaxanthin and a variable concentration of lutein. A further correction to each individual standard was computed by subtraction of the minor contribution (< 5%) of lutein in zeaxanthin (and vice versa) prior to linear regression of the ratios of corrected peak area against concentration. Lutein content in an unknown sample was interpolated from this calibration curve.

2.4. Sample preparation

Sample preparation was adapted from that used for the rapid analysis of carotenoids in bovine milk (Indyk, 1987). Approximately 1 g of powder or 1 g liquid test sample was weighed accurately into a 60 mL boiling tube to which 10 mL of ethanolic pyrogallol (1% w/v) was added. For heterogeneous products, a 4 g aliquot of a slurry (25 g powder in 100 g water) was used. A 1 mL aliquot of zeaxanthin intermediate standard (0.9 mg mL⁻¹) was added to each sample, followed by 2 mL of a 50% w/v KOH

solution, then the tubes were incubated at 70 °C for 10 min in a water bath with periodic agitation to avoid powder agglomeration and ensure efficient lipid saponification. Following cooling in a cold water bath, 20 mL of extraction solution (hexane:diisopropyl ether, 75:25 v/v) was added and the tubes placed on a horizontal shaker for 5 min. Water (20 mL) was added, the tubes capped, inverted 10 times and centrifuged for 10 min at 200 × g. A 1 mL aliquot of the upper solvent layer was transferred to an HPLC vial, blown to dryness, and reconstituted with 1 mL mobile phase ready for analysis.

2.5. Chromatography

The reversed phase column selected for use was a YMC Carotenoid C₃₀, 250 × 4.6 mm, 3 μm (Waters, Milford, MA, USA). The mobile phase consisted of a methanol:dichloromethane (70:30 v/v) run isocratically at a flow rate of 0.5 mL min⁻¹. The injection volume was 50 μL and detection was by photodiode array (370–600 nm) with quantitation at 450 nm.

2.6. Statistical analysis

Statistical evaluation of experimental data for method performance parameters was accomplished with Microsoft Excel 2003 SP1 (Microsoft Corporation, Redmond, WA, USA).

3. Results

3.1. Method performance

Chromatography achieved under the described conditions is illustrated in [Figure 1](#) for carotenoid standards and representative samples. Baseline separation of lutein and zeaxanthin was achieved under the described conditions, and run time was optimised for the polar xanthophylls. Peak identity for the xanthophylls was confirmed by equivalence of retention time and UV–vis spectra against authentic standards.

The performance of the method was determined as summarized in [Table 1](#). Linearity of detector response for both lutein and zeaxanthin was confirmed by least-squares regression analysis of multilevel calibration standards. In the absence of a currently available infant formula certified reference material with certified levels of lutein, method accuracy was confirmed by evaluating spiked recovery at 3 levels equivalent to 10, 20 and 30 mg dL⁻¹ of ready-to-feed infant formula. In addition, accuracy was evaluated for 2 different lutein-supplemented samples by comparing lutein levels as measured by the described procedure against the values obtained from the aggregate of multiple extractions (n = 4). Precision, as repeatability, was determined using 6 replicate analyses of a sample

of a lutein fortified infant formula and this was repeated on 3 separate days to provide an estimate of intermediate precision. The method detection limit (MDL) was determined in accordance with US Environmental Protection Agency procedures (USEPA, 1993).

Recoveries of lutein and zeaxanthin were also quantitated independently through each of the separate saponification and extraction steps (Table 2). Recovery through saponification (70 °C for 10 and 30 min) was determined following acid neutralization after hydrolysis and direct dilution in ethanol without solvent extraction, a modification of a method reported for vitamin A (De Vries & Silvera, 2002). Recovery attributed to the described single solvent extraction procedure was assessed by comparison of peak area of each carotenoid with that of unextracted analyte.

3.2. Milk and milk powder samples

The described method was applied to the analysis of various pediatric formulas, cow and goat whole milk powders, and liquid cow and human milk; results summarised in Table 3. Endogenous lutein levels measured in reconstituted infant formulas and bovine whole milk powders ranged from 1.0 to 5.8 mg dL⁻¹. The levels in fluid bovine and human milk are comparable at 6.0 and 4.3 mg dL⁻¹, respectively.

4. Discussion

It has been recently identified that there is a need for validated methods for the analysis of carotenoids, including lutein, in fortified foods (Blake, 2007). The method presently described was developed predominantly for the compliance testing of lutein supplemented pediatric formulas, while also being suitable for estimation of endogenous lutein in milk. In a high sample throughput laboratory, simple and rapid procedures for extraction of lipophilic components are preferred over those requiring multiple solvent extractions.

Due to the high fat content of milk products, either alkaline hydrolysis or lipase digestion is generally advocated to both remove lipid and release carotenoids (Oliver & Palou, 2000; Schweigert et al., 2000). Although saponification has been employed in the analysis of lutein in milk (Calderón et al., 2007; Jewell et al., 2004; Lietz et al., 2006), the polar xanthophylls are reportedly vulnerable to alkaline conditions and lutein losses have been noted (Liu et al., 1998; Rodríguez-Bernaldo de Quirós & Costa, 2006; Scott, 1992), since apart from degradation via cleavage, *cis*–*trans* isomerisation is also possible. It is therefore expedient to apply the mildest conditions necessary to effectively hydrolyse triglycerides. The impact of high temperature alkaline hydrolysis on lutein and zeaxanthin recovery

was assessed directly, without solvent extraction, and losses due to either mechanism during the described saponification procedure were confirmed as analytically insignificant for both carotenoids.

Hexane-based extraction solvent mixtures containing varying proportions of diisopropyl ether or ethyl acetate were evaluated during extraction of the non-saponifiable carotenoid components. Although the more polar hexane:ethyl acetate system gave predictably higher absolute lutein recoveries, quantitative recovery in a single extraction was not achieved and higher levels of ethyl acetate resulted in degradation of chromatographic performance. In contrast, while the single-extraction procedure incorporating diisopropyl ether was also less than quantitative, the extract yielded unequivocal chromatography. Both binary solvent systems were found to require at least 3 extractions for quantitative recovery. The inability to quantitatively partition lutein and zeaxanthin in a single extraction was not unexpected in view of their relative high polarity compared to the parent carotenes.

In the absence of quantitative recovery during a single-extraction scheme, or in procedures involving multiple steps, the use of an internal standard is mandatory. Since zeaxanthin is structurally comparable to lutein and further, was found in insignificant levels in test samples, it was therefore considered to be an ideal internal standard for the specific determination of lutein by LC-UV techniques.

The utility of zeaxanthin was also supported by baseline resolution from lutein under the described chromatographic conditions, in contrast to several previous reports where aggregate values of lutein and zeaxanthin were a consequence of inadequate resolution (Canfield et al., 2003; Jackson et al., 1998; Liu et al., 1998). Since the solubility of zeaxanthin in ethanol is lower than lutein, it proved necessary to sonicate and filter the stock standard solutions until a stable absorbance (at 450 nm) was achieved prior to calibration.

Recovery and precision were assessed using the criteria developed by AOAC International (2004). The estimated recovery range of 101–108% is within the guideline limits of 75–120% at typical lutein concentration levels found in milk and supplemented formula, while the measured HorRat_r (0.3) is consistent with the guideline range (0.3–1.3).

While LC–MS methods may assist in the identification of carotenoids, most reports have utilised the photodiode-array detector for confirmation of peak identity by spectral comparison against authentic compounds. In the method described presently, this technique was found to be both facile and adequate for the unequivocal identification of lutein and zeaxanthin as both share similar but distinct spectral characteristics.

The method was applied to the analysis of various pediatric formulas and milk powders as well as human and bovine milk. The applicability of this method to each matrix was initially assessed by analysing each sample type in the absence of internal standard to reveal the potential contribution of endogenous zeaxanthin. In goat whole milk powder and pediatric formulas, there were negligible amounts of endogenous zeaxanthin. While human and bovine milk showed trace levels of endogenous zeaxanthin, the quantitation of lutein was not significantly affected given the relatively high level of internal standard added.

In contrast to the carotenes, it has been suggested that lutein may be actively secreted into mammalian milk in view of its specific role during early neonatal macular development (Gossage, Deyhim, Yamini, Douglass, & Moser-Veillon, 2002; Jewell et al., 2004). Human milk reportedly contains several predominant carotenoids including α - and β -carotene, lutein, zeaxanthin, β -cryptoxanthin and lycopene, while bovine milk is dominated by β -carotene and lutein (Calderón et al., 2007; Canfield et al., 2003; Giuliano et al., 1992; Gossage et al., 2002; Jackson et al., 1998; Jewell et al., 2004; Khachik et al., 1997; Liu et al., 1998; Nozière et al., 2006; Schweigert et al., 2000). In the present study, a representative human milk sample yielded lutein levels consistent with the range (0.8–5.7 mg dL⁻¹) of previously reported values (Canfield et al., 2003; Gossage et al., 2002; Jackson et al., 1998; Lietz et al., 2006). Similarly, lutein levels measured in raw and processed consumer bovine milk were also comparable and recent reports (trace–6.6 mg dL⁻¹) (Calderón et al., 2006; Calderón et al., 2007; Indyk, 1987; Nozière et al., 2006). Minimal losses have been reported for lutein through thermal milk processing (Nozière et al., 2006), and indeed, only a small decrease was observed in this study. There are no reported literature values for the lutein content in goat milk, although this species milk is characterised by the absence of β -carotene (Nozière et al., 2006). Prior to the present study, one report only has surveyed xanthophyll content of ready-to-feed formula milk. In that study, variable levels of lutein (trace–18.0 mg dL⁻¹) and zeaxanthin (trace–2.3 mg dL⁻¹) were found, although it was unknown which formulas were intentionally lutein-supplemented, and it was speculated that egg lipid, a commonly used ingredient, may also contribute to the xanthophylls contribution (Jewell et al., 2004). In the present study, 2 formulas were intentionally supplemented with lutein during manufacture, while the remainder contained endogenous levels generally consistent with the contribution of milk fat.

5. Conclusions

Infant formulas are increasingly supplemented with lutein due to its putative role as a protective factor in the neonatal retinal epithelium. A rapid, routine and robust method for the compliance analysis of lutein is therefore essential. The method described incorporates a mild alkaline hydrolysis and single-

extraction technique combined with an isocratic chromatographic separation and has been demonstrated to be applicable to both supplemented milk-based pediatric formulas and endogenous lutein in milk.

Acknowledgements

Support for this work from Fonterra Co-operative Group Ltd. (New Zealand) is gratefully acknowledged.

References

- Aman, R., Bayha, S., Carle, R., & Schieber, A. (2004). Determination of carotenoid stereoisomers in commercial dietary supplements by high-performance liquid chromatography. *Journal of Agricultural and Food Chemistry*, 52, 6086–6090.
- AOAC International. (2004). Definitions and calculations of HORRAT values from intra-laboratory data 2004-01-18. Available from. <http://www.aoac.org>. Accessed 14.02.08.
- Ball, G.F.M. (2006). *Vitamins in foods: analysis, bioavailability and stability*. (pp. 422–427). Boca Raton, FL, USA: CRC Taylor & Francis.
- Barua, A.B., Olson, J.A., Furr, H.C., & van Breemen, R.B. (2000). Vitamin A and carotenoids. In A. P. De Leenheer, W. E. Lambert, & J. F. Bocxlaer (Eds.), *Modern chromatographic analysis of vitamins* (pp. 1–74). New York, NY, USA: Marcel Dekker.
- Blake, C.J. (2007). Status of methodology for the determination of fat-soluble vitamins in foods, dietary supplements and vitamin premixes. *Journal of AOAC International*, 90, 897–910.
- Breithaupt, D.E. (2004). Simultaneous HPLC determination of carotenoids used as food coloring additives: applicability of accelerated solvent extraction. *Food Chemistry*, 86, 449–456.
- Calderón, F., Chauveau-Duriot, B., Martin, B., Graulet, B., Doreau, M., & Nozière, P. (2007). Variations in carotenoids, vitamins A and E, and color in cow's plasma and milk during late pregnancy and the first three months of lactation. *Journal of Dairy Science*, 90, 2335–2346.
- Calderón, F., Tornambé, G., Martin, B., Pradel, P., Chauveau-Duriot, B., & Nozière, P. (2006). Effects of mountain grassland maturity stage and grazing management on carotenoids in sward and cow's milk. *Animal Research*, 55, 533–544.
- Canfield, L.M., Clandinin, M.T., Davies, D.P., Fernandez, M.C., Jackson, J., Hawkes, J., et al. (2003). Multinational study of major breast milk carotenoids of healthy mothers. *European Journal of Nutrition*, 42, 133–141.

- Canfield, L.M., Giuliano, A.R., & Graver, E.J. (1995). Carotenoids, retinoids, and vitamin K in human milk. In R. G. Jensen (Ed.), *Handbook of milk composition* (pp. 693–705). San Diego, CA, USA: Academic Press.
- Craft, N.E., & Soares, J.H., Jr. (1992). Relative solubility, stability, and absorptivity of lutein and β -carotene in organic solvents. *Journal of Agricultural and Food Chemistry*, 40, 431–434.
- Dachtler, M., Glaser, T., Kohler, K., & Albert, K. (2001). Combined HPLC–MS and HPLC–NMR on-line coupling for the separation and determination of lutein and zeaxanthin stereoisomers in spinach and in retina. *Analytical Chemistry*, 73, 667–674.
- De Vries, J.W., & Silvera, K.R. (2002). Determination of vitamins A (retinol) and E (alpha-tocopherol) in foods by liquid chromatography: collaborative study. *Journal of AOAC International*, 85, 424–434.
- Eitenmiller, R.R., & Landen, W.O., Jr. (1999). Vitamin A and carotenoids. In R.R. Eitenmiller, & W.O. Landen, Jr. (Eds.), *Vitamin analysis for the health and food sciences* (pp. 3–75). Boca Raton, FL, USA: CRC Press.
- Giuliano, A.R., Neilson, E.M., Kelly, B.E., & Canfield, L.M. (1992). Simultaneous quantitation and separation of carotenoids and retinol in human milk by high performance liquid chromatography. *Methods in Enzymology*, 213, 391–399.
- Gossage, C.P., Deyhim, M., Yamini, S., Douglass, L.W., & Moser-Veillon, P. B. (2002). Carotenoid composition of human milk during the first month post-partum and the response to β -carotene supplementation. *American Journal of Clinical Nutrition*, 76, 193–197.
- Granelli, K., & Helmersson, S. (1996). Rapid high performance liquid chromatographic method for determination of β -carotene in milk. *Journal of Chromatography A*, 721, 355–358.
- Hart, D.J., & Scott, K.J. (1995). Development and evaluation of an HPLC method for the analysis of carotenoids in foods, and the measurement of the carotenoid content of vegetables and fruits commonly consumed in the UK. *Food Chemistry*, 54, 101–111.
- Indyk, H.E. (1987). The rapid determination of carotenoids in bovine milk using HPLC. *Journal of Micronutrient Analysis*, 3, 169–183.
- Jackson, J.G., Lien, E.L., White, S.J., Bruns, N.J., & Kuhlman, C.F. (1998). Major carotenoids in mature human milk: longitudinal and diurnal patterns. *The Journal of Nutritional Biochemistry*, 9, 2–7.
- Jensen, R.G. (1995). Fat-soluble vitamins in bovine milk. In R.G. Jensen (Ed.), *Handbook of milk composition* (pp. 718–725). San Diego, CA, USA: Academic Press.

- Jewell, V.C., Mayes, C.B.D., Tubman, T.R.J., Northrop-Clewes, C.A., & Thurnham, D.I. (2004). A comparison of lutein and zeaxanthin concentrations in formulas and human milk samples from Northern Ireland mothers. *European Journal of Clinical Nutrition*, 58, 90–97.
- Khachik, F., Spangler, C.J., Smith, J.C., Jr., Canfield, L.M., Steck, A., & Pfander, H. (1997). Identification, quantitation, and relative concentrations of carotenoids and their metabolites in human milk and serum. *Analytical Chemistry*, 69, 1873–1881.
- Konings, E.J.M., & Roomans, H.S. (1997). Evaluation and validation of an LC method for the analysis of carotenoids in vegetables and fruit. *Food Chemistry*, 59, 599–603.
- Larsen, E., & Christensen, L.P. (2005). Simple saponification method for the quantitative determination of carotenoids in green vegetables. *Journal of Agricultural and Food Chemistry*, 53, 6598–6602.
- Li, L., Qin, J., Yin, S.-A., & Tang, G. (2007). Effects of mobile phase ratios on the separation of plant carotenoids by HPLC with a C30 column. *Chromatographia*, 65, 91–94.
- Lietz, G., & Henry, C.J.K. (1997). A modified method to minimise losses of carotenoids and tocopherols during HPLC analysis of red palm oil. *Food Chemistry*, 60, 109–117.
- Lietz, G., Mulokozi, G., Henry, J.C. K., & Tomkins, A.M. (2006). Xanthophyll and hydrocarbon carotenoid patterns differ in plasma and breast milk of women supplemented with red palm oil during pregnancy and lactation. *Journal of Nutrition*, 136, 1821–1827.
- Liu, Y., Xu, M.J., & Canfield, L.M. (1998). Enzymatic hydrolysis, extraction, and quantitation of retinol and major carotenoids in mature human milk. *Nutritional Biochemistry*, 9, 178–183.
- Mercadante, A.Z., Steck, A., & Pfander, H. (1999). Carotenoids from Guava (*Psidium guajava* L): isolation and structure elucidation. *Journal of Agricultural and Food Chemistry*, 47, 145–151.
- Nozière, P., Graulet, B., Lucas, A., Martin, B., Grolier, P., & Doreau, M. (2006). Carotenoids for ruminants: from forages to dairy products. *Animal Feed Science and Technology*, 131, 418–450.
- Oliver, J., & Palou, A. (2000). Chromatographic determination of carotenoids in foods. *Journal of Chromatography A*, 881, 543–555.
- Panfili, G., Fratianni, A., & Irano, M. (2004). Improved normal high performance liquid chromatography procedure for the determination of carotenoids in cereals. *Journal of Agricultural and Food Chemistry*, 52, 6373–6377.
- Rodríguez-Bernaldo de Quirós, A., & Costa, H.S. (2006). Analysis of carotenoids in vegetable and plasma samples: a review. *Journal of Food Composition and Analysis*, 19, 97–111.

- Rousell, R., Raley, L., & Hofsommer, H.J. (1996). Application of diode array detection with a C-30 reversed phase column for the separation and identification of saponified orange juice carotenoids. *Journal of Agricultural and Food Chemistry*, 44, 2176–2181.
- Sauvant, P., Grolier, P., & Azaïs-Braesco, V. (2002). Vitamin A, nutritional significance. In J. W. Fuquay, P. F. Fox, & H. Roginski (Eds.), *Encyclopedia of dairy sciences*, Vol. 4 (pp. 2657–2664). London, England: Academic Press.
- Schlatterer, J., & Breithaupt, D.E. (2006). Xanthophylls in commercial egg yolks: quantification and identification by HPLC and LC–(APCI)MS using a C30 phase. *Journal of Agricultural and Food Chemistry*, 54, 2267–2273.
- Schweigert, F.J., Hurtienne, A., & Bathe, K. (2000). Improved extraction procedure for carotenoids from human milk. *International Journal for Vitamin and Nutrition Research*, 70, 79–83.
- Scott, K.J. (1992). Observations on some of the problems associated with the analysis of carotenoids in foods by HPLC. *Food Chemistry*, 45, 357–364.
- Scott, K.J., Finglas, P.M., Seale, R., Hart, D.J., & de Froimont-Gortz, I. (1996). Interlaboratory studies of HPLC procedures for the analysis of carotenoids in foods. *Food Chemistry*, 57, 85–90.
- Su, Q., Rowley, K.G., & Balazs, N.D.H. (2002). Carotenoids: separation methods applicable to biological samples. *Journal of Chromatography B*, 781, 393–418.
- Trumbo, P.R., & Ellwood, K.C. (2006). Lutein and zeaxanthin intakes and risk of age-related macular degeneration and cataracts: an evaluation using the Food and Drug Administration's evidence-based review system for health claims. *American Journal of Clinical Nutrition*, 84, 971–974.
- USEPA. (1993). *Guidance on evaluation, resolution, and documentation of analytical problems associated with compliance monitoring*. Washington, DC, USA: U.S. Government Printing Office. USEPA 821-B-93-001.

Table 1. Method performance parameters

Analyte	Range (ng mL ⁻¹)	Linear regression	r ²	MDL ^a (mg hg ⁻¹)	RSD _r ^b (%)	RSD _{IR} ^c (%)	Recovery ^d (%)	Recovery ^e (%)
Lutein	15.7–470.9	y = 1414.4x - 2377.3	0.9999	1.4	2.2	3.2	101–108	95
Zeaxanthin	20.6–618.5	y = 1367.7x - 5936.1	0.9997					

^a Method detection limit: determined from n replicates at or near the expected detection limit, MDL = $t_{(n-1, 1-\alpha)} \times sd$, where n = 7 and $\alpha = 0.05$

^b Relative standard deviation repeatability: RSD_r = $sd/mean \times 100$, n = 6

^c Relative standard deviation intermediate reproducibility: RSD_{IR} = $sd/mean \times 100$, n = 18 (6 samples tested on 3 different days)

^d Mean recovery of lutein spiked blank skim-milk samples (3 levels each in triplicate)

^e Mean recovery of 2 independent samples by the described method, expressed as a percentage of total extractable lutein (measured as the aggregate of 4 pooled extractions quantitated against lutein external standard)

Table 2. Recovery of lutein and zeaxanthin through separate saponification and solvent extraction steps

Analyte	Saponification recovery ^a (%)		Extraction recovery ^b (%)
	70 °C, 10 min	70 °C, 30 min	
Lutein	99 ± 3.7	92 ± 2.2	68
Zeaxanthin	102 ± 4.6	99 ± 4.4	69

^a Mean of 5 replicates ± standard deviation

^b Singlicate determinations

Table 3. Lutein levels in various milk samples and infant formulas

Sample	Lutein ^a (mg L ⁻¹)
Whey-based, partially oil-filled pediatric formula	1.0
Whey-based, milk-fat pediatric formula	1.8
Whey-based, oil-filled pediatric formula ^b	15.3
Whey-based, oil-filled pediatric formula ^b	6.2
Milk-based, milk-fat pediatric formula	2.6
Milk-based, oil-filled pediatric formula	2.6
Cow whole milk powder	5.8
Goat whole milk powder	1.1
Raw whole milk	6.0
Processed whole milk	3.3
Human milk	4.3

^a Values are given on a reconstituted basis normalised to 12% solids

^b Lutein-supplemented infant formula




Figure 1. Overlaid chromatograms of milk and powder samples spiked with zeaxanthin internal standard (column: YMC Carotenoid C₃₀, 250 × 4.6 mm, 3 μm; mobile phase methanol:dichloromethane (70:30 v/v); flow rate: 0.5 mL min⁻¹; detection: 450 nm)